

Estimation of acoustic indices with CVs for Northeast Arctic saithe in the Norwegian coastal survey 2003–2017 applying the Sea2Data StoX software

Sigbjørn Mehl, Espen Johnsen, Åsmund Skålevik and Asgeir Aglen



HAVFORSKNINGSINSTITUTTET

Project Report

Report: FISKEN OG HAVET	No. – Year: 1-2018	Date: 23.01.2018	Distribution: Open
Title: Estimation of acoustic indices with CVs for Northeast Arctic saithe in the Norwegian coastal survey 2003–2017 applying the Sea2Data StoX software			Project no.: 82354
Authors: Sigbjørn Mehl, Espen Johnsen, Åsmund Skålevik and Asgeir Aglen			Assignor(s): Ministry of Trade, Industry and Fisheries
			Program: Coastal ecosystems
			Research group: Demersal fish
			Number of pages in total: 19

Summary (Norwegian):

Sea2Data programmet StoX er brukt til å estimere akustiske indeks med CV og lengde og vekt ved alder for nordøstarktisk sei fra det norske kysttoktet Varanger – Stad i 2013 til 2017. Indeksene er sammenlignet med indeks tidligere estimert med programmet BEAM, og forskjeller er i hovedsak forklart ved forskjeller i strata-areal.

Summary (English):

The Sea2Data software StoX was applied to estimate acoustic indices with CVs and length and weight at age for Northeast Arctic saithe from the Norwegian coastal survey Varanger - Stad in 2013 to 2017. The indices are compared to indices previously estimated by the BEAM software, and differences are mainly explained by different strata areas.

Emneord (norsk):

1. Nordøstarktisk sei
2. Akustiske indeks
3. Norskekysten Varanger-Stad
4. StoX programvare

Subject heading (English):

1. Northeast Arctic saithe
2. Acoustic indices
3. Norwegian coast Varanger-Stad
4. StoX software

Content

1	Background	4
2	Material and Methods.....	5
2.1	Survey operation and data sampled	5
2.2	Acoustic measurements	7
2.3	Sampling of catch and use of age-length data.....	10
2.4	Estimation of variance	10
2.5	StoX input, settings and filters	10
3	Results.....	11
3.1	Total echo abundance of saithe	11
3.2	Saithe abundance indices, ratio StoX/BEAM and CVs	12
3.3	Saithe length and weight at age	15
4	Conclusions.....	16
5	References.....	17
6	Appendix 1. Annual survey reports 2003-2016.....	18

1 Background

The Institute of Marine Research (IMR), Bergen, has performed an acoustic survey annually in October-November since 1985 to obtain indices of abundance and estimates of length and weight at age of saithe north of 62°N. The indices are the only fishery-independent data in the assessment of Northeast Arctic saithe in ICES. In autumn 2003 the saithe- and coastal cod surveys were combined. A new survey was designed, with new stratification and smaller strata based on depth and fish distribution in recent years, and with new and more regular transects (Figure 2.2). However, only the part of the new combined survey representing the traditional saithe survey area have so far been included in the estimation of acoustic abundance indices for saithe in order to maintain a uniform time series for tuning of the assessment model.

The new Sea2Data software StoX was applied to estimate acoustic indices with CVs for saithe for the period 2003 to 2017. The main difference between the SAS based BEAM Program (Totland and Godø 2001) used for saithe until 2016 and StoX acoustic abundance estimation is that in BEAM the survey area is divided into rectangles, and for each rectangle an average acoustic density (sA) is calculated, while in StoX transects are defined within each stratum as primary sampling units (PSUs) and used to calculate acoustic density (Jolly and Hampton 1990).

StoX does not use age-length keys (ALK) in the traditional sense with ALKs estimated for large areas. Missing age information is imputed from known age-length data within station. If age information is still missing StoX searches within strata, or lastly within all strata. If no age is available for a length group, the abundance estimate is presented as unknown age. StoX does also allow for uncertainty estimation by bootstrapping the transects and assigned trawl stations.

2 Material and Methods

2.1 Survey operation and data sampled

Table 2.1 presents the vessels participating in the survey in 2003-2017 with some basic trawl information.

Table 2.1. Sea2Data cruise number, start and end data, serial numbers, number of trawl stations for vessel participating in the Norwegian acoustic coastal survey in 2003-2017.

Year	Vessel	Cruice number	Start	End	Serial number		No. trawl stations
					From	To	
2003	Johan Hjort	2003211	1010	1108	81601	81686	228
	Jan Mayen	2003705	1020	1114	81801	81942	
2004	Johan Hjort	2004212	1011	1111	55001	55090	210
	Jan Mayen	2004704	1015	1109	55201	55320	
2005	Johan Hjort	2005212	1011	1107	55001	55061	164
	Jan Mayen	2005704	1013	1111	55201	55303	
2006	Johan Hjort	2006213	1024	1119	55001	55074	250
	Jan Mayen	2006705	1021	1118	55201	55376	
2007	Johan Hjort	2007212	1015	1110	55001	55079	229
	Jan Mayen	2007703	1023	1119	55201	55350	
2008	Johan Hjort	2008210	1028	1207	55001	55133	237
	Jan Mayen	2008705	1118	1120	00001	00011	
	Håkon Mosby	2008623	1125	1216	55558	55650	
2009	Johan Hjort	2009209	1005	1029	55001	55081	198
	Jan Mayen	2009703	0928	1018	55201	55280	
	Jan Mayen	2009704	1108	1112	55281	55291	
	Håkon Mosby	2009623	1201	1206	55401	55426	
2010	Johan Hjort	2010211	0927	1109	55001	55165	177
	Jan Mayen	2010704	1108	1111	55201	55212	
2011	Johan Hjort	2011214	1006	1028	55001	55050	160
	Helmer Hansen	2011722	1016	1112	55301	55398	
	Helmer Hansen	2011723	1114	1115	55399	55410	
2012	Johan Hjort	2012210	1002	1029	55001	55083	160
	Håkon Mosby	2012620	1015	1113	55301	55377	
2013	Johan Hjort	2013210	1003	1026	55001	55066	162
	Håkon Mosby	2013623	1013	1113	55201	55281	
	Helmer Hansen	2013851	1101	1108	55301	55315	
2014	Johan Hjort	2014213	1002	1031	55001	55096	205
	Håkon Mosby	2014621	0926	1023	55201	55296	
	Helmer Hansen	2014011	1031	1101	55551	55563	
2015	Johan Hjort	2015211	1007	1105	55001	55092	195
	Håkon Mosby	2015621	1002	1028	55201	55289	
	Helmer Hansen	2015843	1103	1106	55301	55314	
2016	Johan Hjort	2016210	1001	1030	55001	55098	216
	Håkon Mosby	2016620	1005	1031	55201	55303	
	Helmer Hansen	2016883	1027	1103	55101	55115	
2017	Johan Hjort	2017210	1005	1115	55001	55118	244
	Kristine Bonnevie	2017620	1001	1107	55201	55326	

Table 2.2 gives an account of the age material for demersal species sampled from all trawl hauls and Table 2.3 gives the area used in the saithe acoustic abundance estimates with BEAM and the corresponding default area in the StoX kysttokt_strata.txt used for saithe, by Main Areas. The StoX area used for saithe was on average 90% of the area used in BEAM.

Table 2.2. Number of demersal fish measured for age in the Norwegian acoustic coastal survey 2003-2017.

Year	Saithe	Haddock	Cod	Beaked redfish	Golden redfish	Hake	Halibut	Anglerfish
2003	1589	3135	3390					
2004	1415	3784	2539					
2005	1052	1828	1669					
2006	1241	1963	1583					
2007	1184	2066	1183					
2008	1153	1850	1785	13	262		31	31
2009	883	1402	2341	56	138		35	33
2010	1116	1632	2466	9	93		43	28
2011	1009	1325	2056	14	93		93	6
2012	1078	1500	2196	63	226	170	39	17
2013	905	1408	2432	106	255	44	49	7
2014	1139	1710	2632	129	188	122	39	8
2015	1627	1711	2424	179	544	156	39	12
2016	1448	1895	2931	209	577	114	69	23
2017	1612	2314	2653	212	918	192	95	30

Table 2.3. Area (NM^2) used in the saithe acoustic abundance estimation with BEAM in 2003-2016 and corresponding default area in the StoX kysttokt_strata.txt used for saithe, by Main Areas

Year	A 6930- 7130	B 6700- 6930	C 6330- 6700	D 6200- 6330	Total
2003	8117.6	8893.0	2149.8	3243.8	22404.2
2004	8552.0	8354.3	2546.4	5171.8	24624.4
2005	9224.8	9041.1	3342.5	4115.6	25723.9
2006	8601.8	8938.8	2546.4	4327.6	24414.7
2007	8601.8	9041.1	2745.3	4327.6	24715.8
2008	9224.8	8219.0	3740.4	4327.6	25511.9
2009	8601.8	6876.8	2944.3	4327.6	22750.6
2010	9224.8	7633.5	2944.3	4327.6	24130.3
2011	8414.9	7161.5	3094.3	4327.6	22998.4
2012	8414.9	7735.5	1750.3	4327.6	22228.3
2013	8626.8	6088.5	2944.3	4327.6	21987.3
2014	9254.8	7625.0	3194.3	4327.6	24401.8
2015	8322.8	7335.0	2746.3	4327.6	22731.8
2016	9254.8	7745.0	2746.3	3986.5	23732.6
StoX saithe	8882.7	6979.2	1783.4	4324.7	21970.0

2.2 Acoustic measurements

The method is explained by Dalen and Nakken (1983), MacLennan and Simmonds (1991) and Nedreaas (1997). The acoustic equipment has been continuously improved. Since the early 1990s Simrad EK500 echo sounder and Bergen Echo Integrator (BEI, Knudsen 1990) were used. The Simrad EK60 echo sounder and the Large Scale Survey System (LSSS, Korneliussen *et al.* 2016) replaced the EK500 and BEI; on R/V "Johan Hjort" since the 2005 survey and on R/V "Helmer Hanssen" since the 2008 survey. Since the 2017 survey both R/V "Johan Hjort" and R/V "Kristine Bonnevie" have used the new Simrad EK80 echo sounder.

In the mid-1990s the echo sounder transducers were moved from the hull to a retractable centreboard, on R/V "Johan Hjort" since the 1994 survey and on R/V "Helmer Hanssen" since the 2008 survey. This latter change has largely reduced the signal loss due to air bubbles in the close to surface layer.

The acoustic backscattering values (s_A) are stored at high resolution in LSSS. After scrutinizing and allocating the values to species or species groups, the values are stored with 10 m vertical resolution and 1 nautical mile (NM) horizontal resolution. The procedure for allocation by species is based on:

- composition in trawl catches (pelagic and demersal hauls)
- the appearance of the echo recordings
- inspection of target strength distributions
- inspection of target frequency responses

For each trawl catch the relative s_A -contribution from each species is calculated (Korsbrekke 1996) and used as a guideline for the allocation. If the trawl catch gives the true composition of the species contributing to the observed s_A value, those catch-based s_A - proportions could be used directly for the allocation. In the scrutinizing process, the scientists should evaluate to what extent these catch-based s_A - proportions are reasonable, or if they should be modified based on knowledge about the fish behaviour and the catching performance of the gear.

The survey area is divided into four Main Areas (A, B, C and D, Fig 2.1). Each area is divided into a number of strata, which is defined from polygons where also a "donut variant" exists, i.e. a stratum completely encloses another stratum. Within each stratum, the acoustic course tracks are divided into transects (Fig. 2.2). For the time series 2003-2016 this was done by first running a R-script tagging all the transects as defined in the Excel spreadsheets applied for acoustic abundance estimation of haddock in the coastal survey (Mehl *et al.* 2016) and then the transects are inspected and in some cases edited manually in StoX.

The conversion of mean nautical area scattering coefficient (NASC) ($m^2 \text{ nmi}^{-2}$) to density of fish followed a standard procedure where at least 3 trawl stations (with a catch of more than 2 individuals of saithe) were assigned to each PSU. As a rule, all stations within a stratum was assigned to the PSUs in the same stratum, however, if less than 3 trawl stations had been carried out in a stratum, stations in neighboring strata were assigned to the PSUs such that at least 3 stations were assigned to each PSU.




Figure 2.1. Strata and Main Areas (A,B,C and D) used for acoustic estimations with StoX. The Main Areas were also used for acoustic estimations with BEAM.




Figure 2.2. Standard transects in the new combined saithe and coastal survey.

The combined length distribution (d) was calculated for each transect (PSU (j)) as:

$$d_{l,j} = \sum_{s=1}^s d_{l,s,j}$$

where $d_{l,s,j}$ is density (number by 1 n.mi. tow distance) by 1 cm length group (l) for the stations (s) assigned to PSU (j).

The areal density of fish (ρ) (n per nmi²) by length group l by transect j was calculated as

$$\rho_{j,l} = \frac{\text{NASC}_{j,l}}{\sigma_l}$$

where $\text{NASC}_{j,l}$ is the mean nautical area scattering coefficient by transect (j) and length group (l) and σ_l is the acoustic backscattering cross-section for a fish of length l .

$\text{NASC}_{j,l}$ is calculated as:

$$\text{NASC}_{j,l} = \text{NASC}_j \frac{\sigma_{l,p}}{\sum_l \sigma_{l,p}}$$

where $\sigma_{l,p}$ is the acoustic backscattering cross-section for a fish of length l multiplied with the proportion (p) of a fish of length l in the total length distribution and NASC_j is the mean nautical area scattering coefficient in transect.

The acoustic backscattering cross-section (m²) for a fish of length l is calculated as

$$\sigma_l = 4\pi 10^{(\frac{TS_l}{10})}$$

where the target strength, TS , for a fish of length l (cm) is calculated as

$$TS_l = m \log_{10}(l) + a$$

Where m and a are constants. We applied

$$TS = 20 \log(l) - 68 \quad (\text{Foote, 1987}),$$

The abundance (N) of saithe by length group (l) for stratum k is:

$$N_{k,l} = \rho_{k,l} A_k,$$

where A is stratum area and the mean density of saithe of length group l and stratum k is:

$$\rho_{k,l} = \frac{1}{n_k} \cdot \sum_{j=1}^{n_k} w_{kj} \rho_{kj,l}$$

where $w_{kj} = L_{kj} / \bar{L}_k$ ($j = 1, 2, \dots, n_k$) are the lengths of the n_k sample transects.

Estimates by length are converted to estimates by age using available age-length data from all selected (filtered) stations for the stratum, weighted by station density. The total biomass is estimated by multiplying the numbers at age by weight at age. The abundance by stratum is then summed for defined main areas (Figure 2.1).

2.3 Sampling of catch and use of age-length data

Sorting, weighing, measuring and sampling of the catch are done according to instructions given in Mjanger *et al.* (2017). Since 1999 all data except age are recorded electronically by Scantrol Fishmeter measuring board, connected to stabilized scales. The whole catch or a representative sub sample of most species was length measured on each station. At each trawl station age (otoliths) were sampled from 5 saithe per 5 cm length-group.

2.4 Estimation of variance

The acoustic survey indices of saithe made with StoX are presented together with an estimate of uncertainty (coefficient of variation; CV). These estimates were made using StoX with a stratified bootstrap routine treating each transect as the primary sampling unit. In addition, a bootstrap routine for all trawl stations by strata is carried out by run.

The estimated CV (Standard Deviation · 100/mean) is estimated from 500 iterations and is strongly dependent on the choice of estimator for the indices.

2.5 StoX input, settings and filters

StoX version 2.5 and Rstox 1.7 was used for estimation of acoustic indices and CVs (<http://www.imr.no/forskning/prosjekter/stox/en>). R for Windows version 3.4.3 was used in the R calls (<https://www.r-project.org/>).

Biotic and acoustic XML-files were with a few exceptions downloaded from:
<http://tomcat7.imr.no:8080/DatasetExplorer/v1/html/main.html>.

In **FilterAcoustic**, **FreqExpr** was set to **frequency=38000**. In **NASCExpr**, **acocat** was 22 for saithe.

In **NASC** and **LayerType** was set to **WaterColumn**.

Under **FilterBiotic** and **FishStationExpr**, the following filter were applied:
fs.getLengthSampleCount('SEI') > 2 filtering out stations with less than three specimen (see Johnsen et al. 2016 for more info about filters).

Under **StationLengthDist** and **LengthDistType**, **NormLengthDist** was used, and under **RegroupLengthDist** and **LengthInterval**, **1.0** is applied.

In **DefineStrata**, **kysttokt_strata.txt**. In **StratumArea** and **AreaMethod**, **Accurate** was applied.

Under **BioStationAssignment** and **AssignmentMethod**, **UseProcessData** was used, i.e. assignments from the KT-program with adjustments for 2003-2016. **EstLayers** was set to **1~PELBOT**.

Under **BioStationWeighting** and **WeightingMethod**, **SumWeightedCount** was used.

In **AcusticDensity**, **m** was set to **20** and a to **-68**.

Under **SuperIndAbundance** and **AbundWeightMethod**, **StationDensity** was used, with **LengthDist** set to **RegroupLengthDist**.

3 Results

3.1 Total echo abundance of saithe

Table 3.1 presents the time series of total echo abundance (echo density multiplied by area) of saithe in the investigated areas for StoX and BEAM estimates, and ratio between StoX and BEAM echo abundance estimates.

Table 3.1. Total echo abundance of saithe in Norwegian acoustic coastal survey 2003-2017 (m^2 reflecting surface $\cdot 10^{-3}$) estimated by StoX and BEAM (until 2016), and ratio StoX/BEAM.

Year	StoX	BEAM	Ratio
2003	1625	1231	1.32
2004	1873	2102	0.89
2005	1179	1348	0.87
2006	1160	1270	0.91
2007	1000	1140	0.88
2008	604	832	0.73
2009	950	1098	0.87
2010	765	973	0.79
2011	533	667	0.80
2012	801	899	0.89
2013	606	736	0.82
2014	719	906	0.79
2015	853	941	0.91
2016	975	1091	0.89
2017	839		

The highest and lowest total echo abundance ratio StoX/BEAM was 1.32 and 0.73, while the average ratio over all years was 0.88. The differences are explained by the differences in areas used, in StoX the areas were on average 90% of the areas used in BEAM (Table 2.3). In 2003 the areas used for the southernmost Area D was 1.33 times higher in the StoX estimates than what was originally used in BEAM. The BEAM areas for Area D was adjusted in 2004. In 2008, the year with the lowest ratio, the BEAM areas in area C was about twice the size of the areas used in StoX. There was a better coverage of Area C in 2008, beyond the saithe standard areas, and this was used in the BEAM estimates in 2008 but not in the present StoX estimates.

3.2 Saithe abundance indices, ratio StoX/BEAM and CVs

Table 3.2.1 presents acoustic abundance indices for saithe age groups 1-15+, where 15+ is the sum of indices for age group 15 and older, in 2003 to 2017. Table 3.2.2 gives the ratio between StoX and BEAM indices by age groups 1-10+, total index and total biomass in 2003 to 2016. The highest and lowest single index ratio for age 2-10+ was 2.59 and 0.25, while the highest and lowest average ratio over all age groups in one year was 1.59 and 0.75, and the highest and lowest average ratio for one age group over all years was 1.17 and 0.83. The overall average index ratio was 0.95, the average total index ratio was 0.91 and the average total biomass ratio was 0.91. Except for 2003 and 2008, the StoX indices are in general about 10% lower than the indices estimated by BEAM. As for the total echo abundance, this is explained by the differences in areas used. In StoX the areas were on average 90% of the areas used in BEAM (Table 2.3). In 2003 the area used for Area D was 1.33 times higher in the StoX estimates than what was originally used in BEAM, while in 2008 the BEAM areas in area C was about twice the size of the areas used in StoX.

Table 3.2.3 presents estimates of coefficients of variation (%) for age groups 1-14. Estimates are based on a stratified bootstrap approach with 500 replicates (with transects being primary sampling unit). A CV of 20 % or less could be viewed as acceptable in a traditional stock assessment approach if the indices are unbiased (conditional on a catchability model). Values above this indicate a highly uncertain index with little information regarding year class strength. CVs for age groups 2-5 are at an acceptable level in most years, for age groups 6-7 in less than half of the years while for age group 1 and age groups older than 8 years CVs are above what could be considered as acceptable in all years.

Table 3.2.1. SAITHE. Abundance indices (numbers in millions) from the Norwegian coastal acoustic surveys 2003-2017 estimated by StoX software. + indicates < 0.005.

Year	Age group															Total	Biomass ('000 t)
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15+		
2003	19.8	51.6	128.7	172.1	47.0	8.18	8.03	1.71	1.63	1.17	0.71	0.29	0.16	0.05	0.07	441.3	357.8
2004	0.01	139.8	196.0	114.7	48.2	16.9	4.32	2.23	2.32	0.61	0.90	0.40	0.18	0.06	0.03	526.6	425.4
2005	4.38	18.1	211.5	49.8	16.1	11.9	7.17	1.84	1.11	0.43	0.25	0.25	0	0	0	322.7	262.3
2006	3.46	88.6	42.2	132.1	13.6	4.40	8.91	5.60	2.11	1.07	0.88	0.30	0.26	0.03	0.04	303.5	258.8
2007	2.02	40.3	90.1	25.8	58.3	6.79	4.06	3.83	3.88	0.34	0.67	0.10	0	0	0	236.3	224.2
2008	0.01	50.9	58.2	16.9	8.24	9.85	2.98	0.90	1.37	0.99	0.16	0.25	0	0.03	0	150.8	125.6
2009	0	50.4	97.4	61.5	7.09	4.05	6.11	1.94	1.90	1.38	0.47	0.80	0.19	0	0	233.1	210.0
2010	0.02	7.60	143.0	22.5	17.1	3.95	1.68	3.58	0.43	0.25	0.18	0.30	0.01	0.20	0	200.8	167.1
2011	0	15.2	42.7	59.6	4.61	4.23	1.07	0.81	0.78	0.19	0.03	0.06	0	0	0	129.4	117.7
2012	0.08	68.5	69.0	29.7	18.8	3.48	2.83	0.32	0.58	0.56	0.08	0.05	0	0	0	193.9	148.6
2013	5.02	12.3	77.1	16.5	13.3	11.6	2.19	1.21	0.61	0.39	0.02	+	0.10	0.14	0	140.5	139.1
2014	2.95	28.4	40.1	70.8	8.73	5.62	5.44	1.61	0.55	0.18	0.43	0.10	0	0	0.02	165.0	166.0
2015	0.06	93.5	72.4	22.7	30.1	6.08	4.22	1.85	0.20	0.14	0.07	0.05	0	0	0	231.4	177.6
2016	0.76	72.6	145.7	32.0	10.5	11.2	4.15	2.04	1.46	0.15	0.22	0.12	0.02	0.05	0	281.1	196.0
2017	35.4	23.6	91.1	63.9	13.3	2.76	5.35	2.21	0.62	0.46	0.01	0.02	0.04	0	0.05	238.8	177.2

Table 3.2.2. SAITHE. Ratio new/old acoustic abundance indices and total biomass from the Norwegian coastal acoustic surveys 2003-2016.

Year	Age group										Total	Biomass
	1	2	3	4	5	6	7	8	9	10+		
2003	1.31	1.66	1.47	1.13	1.80	1.32	1.25	1.43	2.33	1.88	1.41	1.41
2004	-	0.92	0.92	0.97	0.98	0.88	0.92	0.75	0.76	0.70	0.93	0.96
2005	4.02	0.81	0.93	0.74	0.79	0.72	0.93	0.84	0.66	1.00	0.88	0.88
2006	0.79	0.90	0.99	0.92	0.70	0.96	1.05	1.00	1.02	0.74	0.93	0.96
2007	0.40	0.89	0.81	0.95	0.95	0.86	0.70	0.93	0.89	1.05	0.88	0.93
2008	0.41	0.92	0.60	0.58	0.60	0.83	0.74	0.84	1.31	0.88	0.70	0.75
2009	-	0.95	0.79	0.79	0.97	0.78	0.91	2.28	2.59	1.69	0.84	0.93
2010	0.08	0.97	0.77	0.73	0.77	0.99	0.88	1.08	1.62	0.65	0.78	0.83
2011	-	1.19	0.91	0.77	0.89	0.74	1.11	0.25	0.39	2.39	0.84	0.80
2012	1.98	0.94	0.69	0.84	0.80	0.92	0.90	1.04	0.79	1.41	0.81	0.86
2013	1.81	0.96	0.68	0.83	1.22	1.04	0.77	0.89	0.80	0.93	0.81	0.86
2014	0.87	1.13	1.00	0.81	0.59	0.65	0.63	0.63	0.50	0.78	0.87	0.81
2015	0.54	1.18	1.00	0.78	0.88	0.81	0.73	0.58	0.59	0.65	1.00	0.92
2016	0.63	1.35	1.08	0.75	0.68	0.70	0.55	0.66	0.70	0.62	1.02	0.88

Table 3.2.3. SAITHE. Estimates of coefficients of variation (%) for acoustic abundance indices from Norwegian coastal acoustic surveys 2003-2017.

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14
2003	0.40	0.23	0.19	0.16	0.15	0.33	0.31	0.23	0.33	0.37	0.50	0.46	0.43	0.97
2004	1.67	0.21	0.19	0.25	0.28	0.28	0.48	0.65	0.32	0.37	0.46	0.47	0.56	0.88
2005	0.54	0.33	0.14	0.21	0.14	0.26	0.28	0.37	0.54	0.80	0.80	0.80	-	-
2006	0.65	0.19	0.33	0.27	0.22	0.31	0.30	0.45	0.40	0.40	0.36	0.58	0.54	0.71
2007	0.35	0.27	0.22	0.19	0.20	0.22	0.36	0.31	0.31	0.37	0.60	0.75	-	-
2008	1.31	0.20	0.20	0.25	0.30	0.17	0.18	0.31	0.33	0.43	0.51	0.41	-	1.08
2009	-	0.39	0.21	0.12	0.28	0.27	0.25	0.39	0.43	0.40	0.55	0.65	0.43	-
2010	1.68	0.32	0.19	0.19	0.20	0.22	0.20	0.27	0.60	0.35	0.75	0.84	1.20	0.76
2011	-	0.23	0.18	0.16	0.24	0.38	0.40	0.48	0.33	1.11	1.04	1.00	-	-
2012	0.68	0.16	0.15	0.18	0.24	0.21	0.34	0.68	0.33	0.60	0.79	1.29	-	-
2013	0.56	0.17	0.12	0.13	0.31	0.19	0.34	0.41	0.42	0.62	1.09	3.11	0.93	0.82
2014	0.73	0.21	0.22	0.24	0.18	0.21	0.18	0.31	0.43	0.56	0.44	0.83	-	-
2015	1.60	0.17	0.16	0.20	0.22	0.26	0.25	0.31	0.30	0.72	0.49	0.58	-	-
2016	2.23	0.17	0.10	0.14	0.17	0.19	0.22	0.30	0.23	0.81	0.84	0.60	0.65	0.58
2017	0.34	0.61	0.13	0.17	0.20	0.34	0.48	0.45	0.39	0.26	0.73	0.94	0.92	-

3.3 Saithe length and weight at age

Tables 3.3.1 and 3.3.2 present the time series of mean length and mean weight at age for age groups 1-14 in the standard saithe area. Age groups with few observations are marked with “+”, while no observations are marked with “-”. Since StoX does not use age-length keys (ALKs) in the traditional sense with an ALK estimated for large areas as done by the BEAM Program, there are differences in length and weight at age for some age groups in some years. However, the overall average ratio StoX/BEAM 2003-2016 was 1.01 for age 1-8 lengths and for age 1-7 weights it was also 1.01.

Table 3.3.1. SAITHE. Length at age in the Norwegian coastal acoustic surveys 2003-2017 estimated by StoX software. + indicates few samples.

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14
2003	26.1	34.0	39.7	44.1	49.0	59.2	63.6	65.9	69.8	75.2	81.8	+	+	+
2004	28.0	32.3	39.9	46.1	53.5	58.2	70.1	76.1	73.4	77.9	76.6	+	+	+
2005	28.0	36.4	39.5	45.6	51.3	59.1	63.0	68.0	70.9	70.4	+	+	-	-
2006	26.2	35.1	40.7	43.4	51.4	57.7	63.9	67.1	70.2	72.4	75.5	+	+	+
2007	27.0	36.2	40.8	46.7	51.0	57.9	65.7	67.8	68.6	72.7	80.9	+	-	-
2008	26.0	36.8	41.6	48.0	52.1	58.4	61.0	67.5	72.3	76.6	+	+	-	+
2009	-	33.8	41.6	47.6	57.9	62.4	66.5	64.7	68.4	74.9	73.0	77.8	+	-
2010	24.2	34.5	38.4	47.1	57.4	61.0	65.0	66.9	68.9	75.8	+	+	+	+
2011	-	36.8	41.7	44.7	56.7	62.8	69.5	65.7	76.0	+	+	+	-	-
2012	29.0	36.4	42.3	47.3	51.6	60.5	66.5	71.8	66.9	79.5	82.9	87.0	-	-
2013	26.0	36.7	41.1	48.7	55.2	60.0	68.8	74.5	75.3	75.4	78.8	+	+	+
2014	24.3	35.8	44.0	46.7	54.8	60.6	61.4	72.3	76.6	80.2	79.3	85.8	-	-
2015	29.3	34.7	41.1	48.8	53.6	60.0	65.8	71.5	+	+	+	+	-	-
2016	28.5	33.2	38.8	47.1	54.1	60.0	67.0	70.5	72.5	81.8	+	+	+	+
2017	25.1	32.6	39.9	45.7	53.5	63.7	69.6	69.6	69.8	73.1	+	+	+	-

Table 3.3.2. SAITHE. Weight at age in the Norwegian coastal acoustic surveys 2003-2017 estimated by StoX software. + indicates few samples.

Year	1	2	3	4	5	6	7	8	9	10	11	12	13	14
2003	169	391	569	828	1144	2104	2482	2923	3543	4348	6017	+	+	+
2004	230	338	622	920	1495	1951	3510	4462	4276	4715	4588	+	+	+
2005	233	500	626	898	1290	2044	2459	2951	3595	3202	+	+	-	-
2006	178	433	682	794	1357	1882	2600	2926	3464	3948	4413	+	+	+
2007	188	471	649	927	1235	1783	2592	2911	2935	3633	4761	+	-	-
2008	193	459	636	986	1274	1846	2140	2817	3633	4243	+	+	-	+
2009	-	376	690	1010	1828	2374	2879	2636	2973	4050	3344	3771	+	-
2010	146	409	556	1016	1814	2227	2624	2851	3116	4363	+	+	+	+
2011	-	503	735	853	1744	2267	3302	2598	4524	+	+	+	-	-
2012	240	456	682	954	1212	1907	2481	3088	2448	4573	4783	4870	-	-
2013	171	481	690	1097	1551	2050	3170	3799	4020	3840	5044	+	+	+
2014	135	445	826	1006	1538	2096	2201	3428	4269	4679	4762	5647	-	-
2015	237	380	624	1042	1361	1955	2674	3390	+	+	+	+	-	-
2016	227	338	518	944	1422	2009	2730	3411	3690	5757	+	+	+	+
2017	142	335	576	882	1477	2511	3165	3277	3246	3576	+	+	+	-

4 Conclusions

The StoX acoustic estimates deviates somewhat from those obtained by the BEAM Program, mainly due to the use of different strata areas (10%) in the two methods.

It is recommended that the present time series of acoustic abundance indices for the period 2003-2017 obtained by StoX become the “official” time series that are used for stock assessment and other purposes. It is further recommended that StoX is used to estimate acoustic indices with CVs from future coastal surveys. At present indices for age groups 3-7 are used in the assessment model. The CV estimates indicates that indices of saithe older than 7 years should not be used for assessment purposes, and it should be investigated how the CVs of age groups 6-7 could be reduce through a better survey design. However, cohort plots show that the survey tracks a year class reasonably well beyond age 7 (Figure 4.1), and the next benchmark assessment for Northeast Arctic saithe should investigate this further.




Figure 4.1. Log abundance curves for each cohort of Northeast Arctic saithe age 3–14 from the Coastal Surveys in 2005–2017.

5 References

- Dalen, J. and Nakken, O. 1983. On the application of the echo integration method. ICES CM 1983/B: 19, 30 pp.
- Foote, K.G. 1987. Fish target strengths for use in echo integrator surveys. Journal of the Acoustical Society of America, 82: 981-987.
- Johnsen, E., Totland, A., Skålevik, Å., Holmin, A. J. and Dingsør, G. 2016. StoX Reference Guide. Version 20161003. Havforskningsinstituttet, Institute of Marine Research. 167 p.
(<http://www.imr.no/forskning/prosjekter/stox/documentation/en>).
- Jolly, G. M., & Hampton, I. (1990). A stratified random transect design for acoustic surveys of fish stocks. *Canadian Journal of Fisheries and Aquatic Sciences*, 47(7), 1282-1291.
- Knudsen, H.P. 1990. The Bergen Echo Integrator: an introduction. - Journal du Conseil International pour l'Exploration de la Mer, 47: 167-174.
- Korneliussen, R. J., Heggelund, Y., Macaulay, G. J., Patel, D., Johnsen, E., & Eliassen, I. K. (2016). Acoustic identification of marine species using a feature library. *Methods in Oceanography*, 17, 187-205.
- Korsbrekke, K. 1996. Brukerveileitung for TOKT312 versjon 6.3. Intern program dokumentasjon., Havforskningsinstituttet, september 1996. 20s. (upubl.).
- MacLennan, D.N. and Simmonds, E.J. 1991. *Fisheries Acoustics*. Chapman Hall, London, England. 336pp.
- Mehl, S, Aglen, A., Berg, E. Dingsør, G. and Korsbrekke, K. 2016. Akustisk mengdemåling av sei, kyst-torsk og hyse Finnmark – Møre hausten 2016. Acoustic abundance of saithe, coastal cod and haddock Finnmark – Møre Autumn 2016. Toktrapport/Havforsknings-instituttet/ISSN 1503-6294, Nr. 15 – 2016. 38s.
- Mjanger, H., Svendsen, B.V., Fotland, Å., Mehl, S., Salthaug, A. 2017. Håndbok for prøvetaking av fisk og krepsdyr. Versjon 4.0. Januar 2017. (In Norwegian).
- Nedreaas, K.H. 1997. Evaluation of the North-East Arctic saithe (*Pollachius virens*) acoustic survey. ICES CM 1997/Y:20.
- Totland, A. and Godø, OR. 2001. BEAM – an interactive GIS application for acoustic abundance estimation. In T. Nishida, P.R. Kailola and C.E. Hollingworth (Eds): Proceedings of the First Symposium on Geographic Information System (GIS) in Fisheries Science. Fishery GIS Research Group. Saitama, Japan.

6 Appendix 1. Annual survey reports 2003-2016

Berg, E., Korsbrekke, K. og Mehl, S. 2003. Akustisk mengdemåling av sei, kysttorsk og ungsild Finnmark – Møre, hausten 2003. Acoustic abundance of saithe, coastal cod and juvenile herring Finnmark – Møre Autumn 2003. Tokrapport/Havforskningsinstituttet/ISSN 1503-6294/Nr. 23 – 2003. 23s.

Berg, E., Korsbrekke, K., Mehl, S. og Nybakk, A. 2004. Akustisk mengdemåling av sei, kysttorsk og ungsild Finnmark – Møre, miljøundersøkingar i fjordar, hausten 2004. Acoustic abundance of saithe, coastal cod and juvenile herring Finnmark – Møre Autumn 2004. Tokrapport/Havforskningsinstituttet/ ISSN 1503-6294/Nr. 22 – 2004. 25s.

Aglen, A., Berg, E., Mehl, S. og Sunnanå, K. 2005. Akustisk mengdemåling av sei, kysttorsk og ungsild Finnmark – Møre hausten 2005. Acoustic abundance of saithe, coastal cod and juvenile herring Finnmark – Møre Autumn 2005. Tokrapport, Havforskningsinstituttet, ISSN 1503-6294, Nr. 23 – 2005. 20s

Aglen, A., Berg, E., Mehl, S. og Sunnanå, K. 2007. Akustisk mengdemåling av sei, kysttorsk og ungsild Finnmark – Møre hausten 2006. Acoustic abundance of saithe, coastal cod and juvenile herring Finnmark – Møre Autumn 2006. Tokrapport, Havforskningsinstituttet, ISSN 1503-6294, Nr. 1 – 2007. 23s

Aglen, A., Berg, E., Mehl, S. og Sunnanå, K. 2008. Akustisk mengdemåling av sei, kysttorsk og ungsild Finnmark – Møre hausten 2007. Acoustic abundance of saithe, coastal cod and juvenile herring Finnmark – Møre Autumn 2007. Tokrapport, Havforskningsinstituttet, ISSN 1503-6294, Nr. 1 – 2008. 24s.

Aglen, A., Berg, E., Mehl, S. og Sunnanå, K. 2009. Akustisk mengdemåling av sei, kysttorsk og hyse Finnmark – Møre hausten 2008. Acoustic abundance of saithe, coastal cod and haddock Finnmark – Møre Autumn 2008. Tokrapport, Havforskningsinstituttet. ISSN 1503-6294/Nr. 5 -2009. 29s.

Mehl, S, Berg, E. og Aglen, A. 2009. Akustisk mengdemåling av sei, kysttorsk og hyse Finnmark – Møre hausten 2009. Acoustic abundance of saithe, coastal cod and haddock Finnmark – Møre Autumn 2009. Tokrapport, Havforskningsinstituttet, ISSN 1503-6294, Nr. 11 – 2009. 29s.

Mehl, S, Aglen, A og Berg, E. 2010. Akustisk mengdemåling av sei, kysttorsk og hyse Finnmark – Møre hausten 2010. Acoustic abundance of saithe, coastal cod and haddock Finnmark – Møre Autumn 2010. Tokrapport/Havforskningsinstituttet/ISSN 1503-6294, Nr. 9 – 2010. 29s.

Mehl, S, Aglen, A, Berg, E. og Korsbrekke 2011. Akustisk mengdemåling av sei, kysttorsk og hyse Finnmark – Møre hausten 2011. Acoustic abundance of saithe, coastal cod and haddock Finnmark – Møre Autumn 2011. Tokrapport/Havforskningsinstituttet/ISSN 1503-6294, Nr. 2 – 2011. 30s.

Mehl, S, Berg, E. Dingsør, G. and Korsbrekke, K. 2012. Akustisk mengdemåling av sei, kysttorsk og hyse Finnmark – Møre hausten 2012. Acoustic abundance of saithe, coastal cod and haddock Finnmark – Møre Autumn 2012. Tokrapport/Havforskningsinstituttet/ISSN 1503-6294, Nr. 5 – 2012. 32s.

Mehl, S, Aglen, A., Berg, E. Dingsør, G. and Korsbrekke, K. 2013. Akustisk mengdemåling av sei, kysttorsk og hyse Finnmark – Møre hausten 2013. Acoustic abundance of saithe, coastal cod and haddock Finnmark – Møre Autumn 2013. Tokrapport/Havforsknings-instituttet/ISSN 1503-6294, Nr. 3 – 2013. 33s.

Mehl, S, Aglen, A., Berg, E. Dingsør, G. and Korsbrekke, K. 2014. Akustisk mengdemåling av sei, kysttorsk og hyse Finnmark – Møre hausten 2014. Acoustic abundance of saithe, coastal cod and haddock Finnmark–Møre Autumn 2014. Tokrapport/Havforskningsinstituttet/ISSN 1503-6294, Nr. 1 – 2014. 38s.

Mehl, S, Aglen, A., Berg, E. Dingsør, G. and Korsbrekke, K. 2015. Akustisk mengdemåling av sei, kysttorsk og hyse Finnmark – Møre hausten 2015. Acoustic abundance of saithe, coastal cod and haddock Finnmark–Møre Autumn 2015 Tokrapport/Havforskningsinstituttet/ISSN 1503-6294, Nr. 4 – 2015. 38s.

Mehl, S, Aglen, A., Berg, E. Dingsør, G. and Korsbrekke, K. 2016. Akustisk mengdemåling av sei, kyst-torsk og hyse Finnmark – Møre hausten 2016. Acoustic abundance of saithe, coastal cod and haddock Finnmark – Møre Autumn 2016. Tokrapport/Havforsknings-instituttet/ISSN 1503-6294, Nr. 15 – 2016. 38s.

Retur: Havforskningsinstituttet, Postboks 1870 Nordnes, NO-5817 Bergen

HAVFORSKNINGSINSTITUTTET
Institute of Marine Research

Nordnesgaten 50 – Postboks 1870 Nordnes
NO-5817 Bergen
Tlf.: +47 55 23 85 00 – Faks: +47 55 23 85 31
E-post: post@imr.no

www.hi.no